

Afghanistan

Medical Summary

The health risk information presented here is summarized from Shoreland Travax[®], a decision-support tool used by health care providers to perform a detailed health risk analysis based on specific locations, individual travel styles, and traveler risk behaviors. Travax provides practitioners current, independently researched malaria risk and prevention recommendations in a map-based format that goes beyond the annual WHO and US CDC statements included here. Not included here are current reports from Travax of disease outbreaks or environmental events that may pose elevated risks to travelers' health and safety. The Providers section of this site offers a directory of health care providers who utilize Shoreland Travax for travel health counseling. Learn more about the detailed reports and maps available from these practitioners (includes links to samples).

General Information

Afghanistan is a developing nation classified as low income. Located in southern Asia (north and west of Pakistan and east of Iran), the climate is classified as dry (arid) to dry (semi arid), with cooler temperatures in some high altitude areas.

Vaccinations

Routine vaccinations are essential due to a persistent rise of vaccine-preventable diseases (especially markedly high rates of diphtheria, pertussis, and measles) globally. Prior to travel, travelers should be up-to-date with the age-appropriate routine vaccinations recommended by their home country, which may include: COVID-19; *H. influenzae* type B (Hib); hepatitis A; hepatitis B; herpes zoster; human papillomavirus; influenza; measles, mumps, rubella; meningococcal; pneumococcal; polio; rotavirus; tetanus, diphtheria, pertussis (Tdap preferred; consider an early pertussis booster for high-risk travelers); varicella.

Depending on your itinerary, your personal risk factors, and the length of your visit, your health care provider may offer you vaccination against chikungunya, cholera, dengue, rabies, tick-borne encephalitis, typhoid fever, or a one time polio booster if you haven't previously received one for travel.

Malaria

The following is current information as reported by the World Health Organization (WHO) and the US Centers for Disease Control (CDC):

WHO—*International Travel and Health* (current online update, Country List)

(2022) Malaria risk due to *P. falciparum* and *P. vivax* exists from May through November below 2000 m.

- Recommended prevention in risk areas: **C** – Risk of *P. falciparum* malaria. Mosquito bite prevention plus atovaquone–proguanil or doxycycline or mefloquine Mosquito-bite prevention plus atovaquone-proguanil or doxycycline or mefloquine chemoprophylaxis (select according to drug-resistance pattern, reported side-effects and contraindications).^{a, b}

^aAlternatively, for travel to rural areas with low risk of malaria infection, mosquito bite prevention can be combined with stand-by emergency treatment (SBET).

^bIn certain areas with multidrug-resistant malaria, mefloquine chemoprophylaxis is no longer recommended. At present, these areas include Cambodia, south-eastern Myanmar and Thailand.

WHO Country List footnote: When available, the date of the most recent update or confirmation is indicated in parentheses in the country list. If no accurate date is indicated, the most recent update or confirmation was provided prior to 2013.

CDC—*Health Information for International Travel* (current online edition)

Transmission areas

- All areas <2,500 m (~8,200 ft) elevation (April–December)

Drug resistance²

- Chloroquine

Species

- *P. vivax* (primarily)
- *P. falciparum* (less commonly)

Recommended chemoprophylaxis

- Atovaquone-proguanil, doxycycline, mefloquine, tafenoquine.³

² Refers to *P. falciparum* malaria, unless otherwise noted.

³ Tafenoquine can cause potentially life-threatening hemolysis in people with glucose-6-phosphate-dehydrogenase (G6PD) deficiency. Rule out G6PD deficiency with a quantitative laboratory test before prescribing tafenoquine to patients.

Other Concerns

Travelers' Diarrhea

High risk exists throughout the country, including in deluxe accommodations. Community sanitation and food safety measures are generally inadequate. Some itineraries (e.g., remote destinations, austere accommodations) and activities (e.g., ecotourism, eating street or local-market food) further increase risk.

Travelers should observe food and beverage precautions, which reduce the likelihood of illness.

Travelers should carry loperamide for self-treatment of diarrhea and, if risk is moderate to high, an antibiotic to add if diarrhea is severe. Consult a knowledgeable health care provider regarding which antibiotic is appropriate for you and most effective for your destination.

Other Food-Borne Illnesses

Precautions to prevent brucellosis may be needed.

Insect- and Arthropod-Borne Diseases

Leishmaniasis, West Nile virus may pose a risk. Personal protective measures are important.

Other Disease and Health Risks

Additional concerns include anthrax disease, Crimean-Congo hemorrhagic fever, helminths, leptospirosis, plague, sexually transmitted infections, snakebites, tuberculosis.

Safety and Security

Key Safety Risks

- Road conditions and traffic collisions
- Domestic and/or international air travel
- Heightened crime risk for women
- Heightened crime risk for LGBTQ+ travelers
- Arbitrary arrest/unlawful detention
- Natural disasters

Key Security Threats

- Armed conflict
- Terrorist attacks by domestic and/or transnational groups, which may target foreign nationals and/or airports
- Risk of violent civil unrest
- Risk of cross-border threats
- Explosions in public places
- Landmines/unexploded ordnance
- Violent crime
- Kidnapping/hostage taking, which may target foreign nationals
- Random acts of armed violence

Emergency Contacts

The emergency number in major cities is 119; the line may not be answered and local operators do not speak English.

© 2025 Shoreland, Inc. All rights reserved.