

Bolivia

Medical Summary

The health risk information presented here is summarized from Shoreland Travax[®], a decision-support tool used by health care providers to perform a detailed health risk analysis based on specific locations, individual travel styles, and traveler risk behaviors. Travax provides practitioners current, independently researched malaria risk and prevention recommendations in a map-based format that goes beyond the annual WHO and US CDC statements included here. Not included here are current reports from Travax of disease outbreaks or environmental events that may pose elevated risks to travelers' health and safety. The Providers section of this site offers a directory of health care providers who utilize Shoreland Travax for travel health counseling. Learn more about the detailed reports and maps available from these practitioners (includes links to samples).

General Information

Bolivia is a developing nation classified as lower middle income. Located in central South America (south of Brazil and north of Paraguay), the climate classifications range from dry summer to humid equatorial (long dry season), with cooler temperatures in some high-altitude areas.

Vaccinations

Routine vaccinations are essential due to a persistent rise of vaccine-preventable diseases (especially markedly high rates of diphtheria, pertussis, and measles) globally. Prior to travel, travelers should be up-to-date with the age-appropriate routine vaccinations recommended by their home country, which may include: COVID-19; *H. influenzae* type B (Hib); hepatitis A; hepatitis B; herpes zoster; human papillomavirus; influenza; measles, mumps, rubella; meningococcal; pneumococcal; polio; rotavirus; tetanus, diphtheria, pertussis (Tdap preferred; consider an early pertussis booster for high-risk travelers); varicella.

Yellow Fever

An official yellow fever vaccination certificate may be required depending on your itinerary. Vaccination is usually recommended if you'll be traveling in areas where there is risk of yellow fever transmission.

- *Requirement:* A certificate proving yellow fever vaccination is required for travelers aged ≥ 1 year coming from countries with risk of YF transmission. This does not apply to airport transit stops (no exit through immigration checkpoint) in risk countries.
- *Official Status:* listed by WHO as a country where YF transmission risk is present.

Other Vaccines

Depending on your itinerary, your personal risk factors, and the length of your visit, your health care provider may offer you vaccination against chikungunya, mpox, rabies, or typhoid fever.

Malaria

The following is current information as reported by the World Health Organization (WHO) and the US Centers for Disease Control (CDC):

WHO—*International Travel and Health* (current online update, Country List)

(2021) Malaria risk due almost exclusively to *P. vivax* (99.9%) exists throughout the year in the entire country below 2500 m. The risk of malaria is highest in the northern departments of Beni and Pando, especially in the localities of Riberalta, Guayaramerín and Sena.

- Recommended prevention in risk areas: **B** - Risk of *P. vivax* malaria only. Mosquito bite prevention plus chloroquine chemoprophylaxis.^a

^aAlternatively, for travel to rural areas with low risk of malaria infection, mosquito bite prevention can be combined with stand-by emergency treatment (SBET).

WHO Country List footnote: When available, the date of the most recent update or confirmation is indicated in parentheses in the country list. If no date is indicated, the most recent update or confirmation was provided before 2013.

CDC—Health Information for International Travel (current online edition)

Areas with malaria: All areas < 2,500 m (8,202 ft). None in the city of La Paz (see Map 3-18).

- Drug resistance³ : Chloroquine.
- Malaria species: *P. vivax* 93%, *P. falciparum* 7%.
- Recommended chemoprophylaxis: Atovaquone-proguanil, doxycycline, mefloquine, primaquine,⁴ or tafenoquine.⁴

³ Refers to *P. falciparum* malaria unless otherwise noted.

⁴ Primaquine and tafenoquine can cause hemolytic anemia in people with G6PD deficiency. Patients must be screened for G6PD deficiency before starting primaquine or tafenoquine. See *Tafenoquine Approved for Malaria Prophylaxis and Treatment* for more information.

Other Concerns

Travelers' Diarrhea

High risk exists throughout the country, with moderate risk in deluxe accommodations. Community sanitation and food safety measures are generally inadequate. Some itineraries (e.g., remote destinations, austere accommodations) and activities (e.g., ecotourism, eating street or local-market food) further increase risk.

Travelers should observe food and beverage precautions, which reduce the likelihood of illness.

Travelers should carry loperamide for self-treatment of diarrhea and, if risk is moderate to high, an antibiotic to add if diarrhea is severe. Consult a knowledgeable health care provider regarding which antibiotic is appropriate for you and most effective for your destination.

Other Food-Borne Illnesses

Precautions to prevent brucellosis may be needed.

Insect- and Arthropod-Borne Diseases

Chagas' disease (American trypanosomiasis), dengue, leishmaniasis, mayaro virus, West Nile virus, Zika may pose a risk. Personal protective measures are important.

Other Disease and Health Risks

Additional concerns include air pollution, altitude illness, anthrax disease, hantavirus, helminths, leptospirosis, plague, schistosomiasis, sexually transmitted infections, snakebites, tuberculosis.

Safety and Security

Key Safety Risks

- Road conditions and traffic collisions
- Public transportation
- Petty crime
- Heightened crime risk for women
- Food and beverage spiking
- Theft of vehicles
- Scams
- Natural disasters

Key Security Threats

- Risk of violent civil unrest
- Risk of violent protests
- Violent crime

Emergency Phone Number

The police emergency number is 110. The English-speaking tourist police number is [+591] 222-5016.

© 2024 Shoreland, Inc. All rights reserved.