

Haiti

Medical Summary

The health risk information presented here is summarized from Shoreland Travax[®], a decision-support tool used by health care providers to perform a detailed health risk analysis based on specific locations, individual travel styles, and traveler risk behaviors. Travax provides practitioners current, independently researched malaria risk and prevention recommendations in a map-based format that goes beyond the annual WHO and US CDC statements included here. Not included here are current reports from Travax of disease outbreaks or environmental events that may pose elevated risks to travelers' health and safety. The Providers section of this site offers a directory of health care providers who utilize Shoreland Travax for travel health counseling. Learn more about the detailed reports and maps available from these practitioners (includes links to samples).

General Information

Haiti is a developing nation classified as low income. Located in the Caribbean Sea (west of Dominican Republic), the climate is classified as humid equatorial (long dry season).

Vaccinations

Routine vaccinations are essential due to a persistent rise of vaccine-preventable diseases (especially markedly high rates of diphtheria, pertussis, and measles) globally. Prior to travel, travelers should be up-to-date with the age-appropriate routine vaccinations recommended by their home country, which may include: COVID-19; *H. influenzae* type B (Hib); hepatitis A; hepatitis B; herpes zoster; human papillomavirus; influenza; measles, mumps, rubella; meningococcal; pneumococcal; polio; rotavirus; tetanus, diphtheria, pertussis (Tdap preferred; consider an early pertussis booster for high-risk travelers); varicella.

Yellow Fever

Although yellow fever does not occur in Haiti, an official yellow fever vaccination certificate may be required depending on your itinerary.

- *Requirement:* A certificate proving yellow fever vaccination is required for travelers aged ≥ 1 year coming from countries with risk of YF transmission. This does not apply to airport transit stops (no exit through immigration checkpoint) in risk countries.

Other Vaccines

Depending on your itinerary, your personal risk factors, and the length of your visit, your health care provider may offer you vaccination against chikungunya, cholera, dengue, rabies, or typhoid fever.

Malaria

The following is current information as reported by the World Health Organization (WHO) and the US Centers for Disease Control (CDC):

WHO—*International Travel and Health* (current online update, Country List)

(2022) Malaria risk due exclusively to *P. falciparum* exists throughout the year in the entire country, but especially in the southern departments such as Grand'Anse, Nippes, and Sud-Est. No chloroquine-resistant *P. falciparum* has been reported.

- Recommended prevention: **C** – Risk of *P. falciparum* malaria. Mosquito-bite prevention plus atovaquone-proguanil or doxycycline or mefloquine chemoprophylaxis (select according to drug-resistance pattern, reported side-effects and contraindications).^{a, b}

^aAlternatively, for travel to rural areas with low risk of malaria infection, mosquito bite prevention can be combined with stand-by emergency treatment (SBET).

^bIn certain areas with multidrug-resistant malaria, mefloquine chemoprophylaxis is no longer recommended. At present, these areas include Cambodia, south-eastern Myanmar and Thailand.

WHO Country List footnote: When available, the date of the most recent update or confirmation is indicated in parentheses in the country list. If no accurate date is indicated, the most recent update or confirmation was provided prior to 2013.

CDC—Health Information for International Travel (current online edition)

Transmission areas

- All (including Labadee, also known as Port Labadee)

Drug resistance²

- None

Species

- *P. falciparum* (99%)
- *P. malariae* (rare)

Recommended chemoprophylaxis

- Atovaquone-proguanil, chloroquine, doxycycline, mefloquine, tafenoquine³

² Refers to *P. falciparum* malaria, unless otherwise noted.

³ Tafenoquine can cause potentially life-threatening hemolysis in people with glucose-6-phosphate-dehydrogenase (G6PD) deficiency. Rule out G6PD deficiency with a quantitative laboratory test before prescribing tafenoquine to patients.

Other Concerns

Travelers' Diarrhea

High risk exists throughout the country, including in deluxe accommodations. Community sanitation and food safety measures are generally inadequate. Some itineraries (e.g., remote destinations, austere accommodations) and activities (e.g., ecotourism, eating street or local-market food) further increase risk.

Travelers should observe food and beverage precautions, which reduce the likelihood of illness.

Travelers should carry loperamide for self-treatment of diarrhea and, if risk is moderate to high, an antibiotic to add if diarrhea is severe. Consult a knowledgeable health care provider regarding which antibiotic is appropriate for you and most effective for your destination.

Other Food-Borne Illnesses

Precautions to prevent brucellosis, seafood poisoning may be needed.

Insect- and Arthropod-Borne Diseases

Mayaro virus, West Nile virus, Zika may pose a risk. Personal protective measures are important.

Other Disease and Health Risks

Additional concerns include anthrax disease, helminths, leptospirosis, marine hazards, sexually transmitted infections, tuberculosis.

Safety and Security

Key Safety Risks

- Road conditions
- Public transportation
- Domestic and/or international air travel
- Petty crime
- Heightened crime risk for women
- Heightened crime risk for LGBTQ+ travelers
- Theft of vehicles
- Natural disasters

Key Security Threats

- Risk of violent civil unrest
- Risk of violent protests
- Cross-border threats in areas bordering the Dominican Republic
- Violent crime
- Kidnapping, which may target foreign nationals
- Random acts of armed violence

Emergency Contacts

The national emergency number is 114; emergency services may be unreliable.

© 2025 Shoreland, Inc. All rights reserved.