

# Namibia

## Medical Summary

*The health risk information presented here is summarized from Shoreland Travax<sup>®</sup>, a decision-support tool used by health care providers to perform a detailed health risk analysis based on specific locations, individual travel styles, and traveler risk behaviors. Travax provides practitioners current, independently researched malaria risk and prevention recommendations in a map-based format that goes beyond the annual WHO and US CDC statements included here. Not included here are current reports from Travax of disease outbreaks or environmental events that may pose elevated risks to travelers' health and safety. The Providers section of this site offers a directory of health care providers who utilize Shoreland Travax for travel health counseling. Learn more about the detailed reports and maps available from these practitioners (includes links to samples).*

## General Information

Namibia is a developing nation classified as upper middle income. Located in southwest Africa along the Atlantic Ocean, the climate is classified as dry (arid) in the south and along the coast and dry (semi arid) in the north.

## Vaccinations

**Routine vaccinations** are essential due to a persistent global rise of vaccine-preventable diseases (especially markedly high rates of diphtheria, pertussis, and measles). Prior to travel, travelers should be up-to-date with the age-appropriate routine vaccinations recommended by their home country, which may include: COVID-19; *H. influenzae* type B (Hib); hepatitis A; hepatitis B; herpes zoster; human papillomavirus; influenza; measles, mumps, rubella (a single early dose is recommended for travelers aged 6-11 months); meningococcal; pneumococcal; polio; rotavirus; tetanus, diphtheria, pertussis (Tdap preferred; consider an early pertussis booster for high-risk travelers); varicella.

## Yellow Fever

Although yellow fever does not occur in Namibia, an official yellow fever vaccination certificate may be required depending on your itinerary.

- **Requirement:** A certificate proving yellow fever vaccination is required for travelers aged  $\geq 9$  months coming from countries with risk of YF transmission. This also applies to airport transit stops (no exit through immigration checkpoint) longer than 12 hours in risk countries.

## Other Vaccines

Depending on your itinerary, your personal risk factors, and the length of your visit, your health care provider may offer you vaccination against dengue, rabies, or typhoid fever.

## Malaria

The following is current information as reported by the World Health Organization (WHO) and the US Centers for Disease Control (CDC):

### WHO—*International Travel and Health* (current online update, Country List)

(2018) Malaria risk due to *P. falciparum* exists from November through June in the following regions: Ohangwena, Omaheke, Omusati, Oshana, Oshikoto and Otjozondjupa. Risk exists throughout the year along the Kunene river in Kunene Region, the Zambezi river in Zambezi Region, and the Okavango river in Kavango regions (West and East).

- Recommended prevention in risk areas: **C** – Risk of *P. falciparum* malaria. Mosquito-bite prevention plus atovaquone-proguanil or doxycycline or mefloquine chemoprophylaxis (select according to drug-resistance pattern, reported side-effects and contraindications).<sup>a, b</sup>

<sup>a</sup>Alternatively, for travel to rural areas with low risk of malaria infection, mosquito bite prevention can be combined with stand-by emergency treatment (SBET).

<sup>b</sup>In certain areas with multidrug-resistant malaria, mefloquine chemoprophylaxis is no longer recommended. At present, these areas include Cambodia, south-eastern Myanmar and Thailand.

*WHO Country List footnote:* When available, the date of the most recent update or confirmation is indicated in parentheses in the country list. If no accurate date is indicated, the most recent update or confirmation was provided prior to 2013.

## CDC—Health Information for International Travel (current online edition)

### Transmission areas

- In the regions of Kavango (East and West), Kunene, Ohangwena, Omaheke, Omusati, Oshana, Oshikoto, Otjozondjupa, and Zambezi
- Rare in other parts of the country
- No malaria transmission in Windhoek (the capital)

### Drug resistance<sup>1</sup>

- Chloroquine

### Species

- *P. falciparum* (primarily)
- *P. malariae*, *P. ovale*, and *P. vivax* (less commonly)

### Recommended chemoprophylaxis

- Kavango (East and West), Kunene, Ohangwena, Omaheke, Omusati, Oshana, Oshikoto, Otjozondjupa, and Zambezi: Atovaquone-proguanil, doxycycline, mefloquine, tafenoquine<sup>2</sup>
- All other areas with malaria transmission: No chemoprophylaxis recommended (insect bite precautions and mosquito avoidance only)<sup>4</sup>

<sup>1</sup> Refers to *P. falciparum* malaria, unless otherwise noted.

<sup>2</sup> Tafenoquine can cause potentially life-threatening hemolysis in people with glucose-6-phosphate-dehydrogenase (G6PD) deficiency. Test for G6PD deficiency with a quantitative laboratory test before prescribing tafenoquine to patients.

<sup>4</sup> Mosquito avoidance includes applying topical mosquito repellent, sleeping under an insecticide-treated mosquito net, and wearing protective clothing (e.g., long pants and socks, long-sleeve shirt). For additional details on insect bite precautions, see Mosquitoes, Ticks, and Other Arthropods chapter.

## Other Concerns

### Travelers' Diarrhea

High risk exists throughout the country, with moderate risk in deluxe accommodations. Community sanitation and food safety measures are generally inadequate. Some itineraries (e.g., remote destinations, austere accommodations) and activities (e.g., ecotourism, eating street or local-market food) further increase risk.

Travelers should observe food and beverage precautions, which reduce the likelihood of illness.

Travelers should carry loperamide for self-treatment of diarrhea and, if risk is moderate to high, an antibiotic to add if diarrhea is severe. Consult a knowledgeable health care provider regarding which antibiotic is appropriate for you and most effective for your destination.

### Other Food-Borne Illnesses

Precautions to prevent brucellosis may be needed.

### Insect- and Arthropod-Borne Diseases

African trypanosomiasis, tick-bite fever, West Nile virus may pose a risk. Personal protective measures are important.

### Other Disease and Health Risks

Additional concerns include anthrax disease, Crimean-Congo hemorrhagic fever, helminths, leptospirosis, Marburg virus disease, marine hazards, plague, schistosomiasis, sexually transmitted infections, tuberculosis.

## Safety and Security

### Key Safety Risks

- Road conditions and traffic collisions
- Public transportation
- Petty crime
- Heightened crime risk for women
- Theft of vehicles
- Scams

### Key Security Threats

- Cross-border threats in areas bordering Angola
- Landmines
- Occasional violent crime

### Emergency Contacts

The national emergency numbers are 112 and 10111.

---

© 2025 Shoreland, Inc. All rights reserved.