

# Paraguay

## Medical Summary

The health risk information presented here is summarized from Shoreland Travax®, a decision-support tool used by health care providers to perform a detailed health risk analysis based on specific locations, individual travel styles, and traveler risk behaviors. Travax provides practitioners current, independently researched malaria risk and prevention recommendations in a map-based format that goes beyond the annual WHO and US CDC statements included here. Not included here are current reports from Travax of disease outbreaks or environmental events that may pose elevated risks to travelers' health and safety. The Providers section of this site offers a directory of health care providers who utilize Shoreland Travax for travel health counseling. Learn more about the detailed reports and maps available from these practitioners (includes links to samples).

### General Information

Paraguay is a developing nation classified as upper middle income. Located in South America (north of Argentina and south of Brazil), the climate is classified as humid equatorial (long dry season) in central areas and subtropical dry winter in the rest of the country.

### Vaccinations

**Routine vaccinations** are essential due to a persistent global rise of vaccine-preventable diseases (especially markedly high rates of diphtheria, pertussis, and measles). Prior to travel, travelers should be up-to-date with the age-appropriate and risk-based routine vaccinations recommended by their home country, which may include:

- COVID-19
- *H. influenzae* type B (Hib)
- Hepatitis A
- Hepatitis B: Protection is especially important for those at increased risk.
- Herpes zoster
- Human papillomavirus
- Influenza
- Measles, mumps, rubella: A single early dose is recommended for travelers aged 6-11 months.
- Meningococcal
- Pneumococcal
- Polio
- Rotavirus
- Tetanus, diphtheria, pertussis: Tdap preferred; consider an early pertussis booster for high-risk travelers.
- Varicella

### Yellow Fever

An official yellow fever vaccination certificate may be required depending on your itinerary. Vaccination is usually recommended if you'll be traveling in areas where there is risk of yellow fever transmission.

- *Requirement:* Per Paraguay's Ministry of Health, a certificate proving yellow fever (YF) vaccination is required for travelers aged 1-59 years coming from risk zones in Bolivia (Santa Cruz Department), Brazil (Amazonas, Minas Gerais, São Paulo, Rio Grande do Sul, and Roraima states), Colombia (Caquetá, Huila, Nariño, Putumayo, Vaupés, and Tolima departments), Guyana, and Peru (Huánuco, Junín, Madre de Dios, San Martín, and Ucayali regions). This also applies to airport transit stops (no exit through immigration checkpoint) longer than 24 hours in risk zones of these countries. Paraguayan authorities also enforce proof of YF vaccination for citizens and foreign residents traveling to risk zones in the aforementioned countries.
- *Official Status:* listed by WHO as a country where YF transmission risk is present.

### Other Vaccines

Depending on your itinerary, your personal risk factors, and the length of your visit, your health care provider may offer you vaccination against chikungunya, dengue, mpox, rabies, or typhoid fever.

## Malaria

The following is current information as reported by the World Health Organization (WHO) and the US Centers for Disease Control (CDC):

### **WHO—*International Travel and Health* (current online update, Country List)**

(2022) Country certified malaria-free by WHO in 2018.

### **CDC—*Health Information for International Travel* (current online edition)**

No malaria transmission.

## Other Concerns

### Travelers' Diarrhea

High risk exists throughout the country, including in deluxe accommodations. Community sanitation and food safety measures are generally inadequate. Some itineraries (e.g., remote destinations, austere accommodations) and activities (e.g., ecotourism, eating street or local-market food) further increase risk.

Travelers should observe food and beverage precautions, which reduce the likelihood of illness.

Travelers should carry loperamide for self-treatment of diarrhea and, if risk is moderate to high, an antibiotic to add if diarrhea is severe. Consult a knowledgeable health care provider regarding which antibiotic is appropriate for you and most effective for your destination.

### Other Food-Borne Illnesses

Precautions to prevent brucellosis may be needed.

### Insect- and Arthropod-Borne Diseases

Chagas' disease (American trypanosomiasis), leishmaniasis, Zika may pose a risk. Personal protective measures are important.

## Other Disease and Health Risks

Additional concerns include anthrax disease, hantavirus, helminths, leptospirosis, New World screwworm (myiasis), plague, sexually transmitted infections, snakebites, tuberculosis.

## Safety and Security

### Key Safety Risks

- Road conditions and traffic collisions
- Public transportation
- Maritime safety
- Petty crime
- Food and beverage spiking
- Theft of vehicles
- Scams
- Natural disasters

### Key Security Threats

- Risk of cross-border threats in areas bordering Argentine, Bolivia, and Brazil
- Occasional violent crime
- Random acts of armed violence

## Emergency Contacts

The police emergency number is 911.

---

© 2026 Shoreland, Inc. All rights reserved.