

Vanuatu

Medical Summary

The health risk information presented here is summarized from Shoreland Travax®, a decision-support tool used by health care providers to perform a detailed health risk analysis based on specific locations, individual travel styles, and traveler risk behaviors. Travax provides practitioners current, independently researched malaria risk and prevention recommendations in a map-based format that goes beyond the annual WHO and US CDC statements included here. Not included here are current reports from Travax of disease outbreaks or environmental events that may pose elevated risks to travelers' health and safety. The Providers section of this site offers a directory of health care providers who utilize Shoreland Travax for travel health counseling. Learn more about the detailed reports and maps available from these practitioners (includes links to samples).

General Information

Vanuatu is a developing nation classified as lower middle income. A chain of islands located in the Pacific Ocean (west of Fiji and north of New Caledonia), the climate is classified as humid equatorial (no dry season).

Vaccinations

Routine vaccinations are essential due to a persistent global rise of vaccine-preventable diseases (especially markedly high rates of diphtheria, pertussis, and measles). Prior to travel, travelers should be up-to-date with the age-appropriate and risk-based routine vaccinations recommended by their home country, which may include:

- COVID-19
- *H. influenzae* type B (Hib)
- Hepatitis A
- Hepatitis B: Protection is especially important for those at increased risk.
- Herpes zoster
- Human papillomavirus
- Influenza
- Measles, mumps, rubella: A single early dose is recommended for travelers aged 6-11 months.
- Meningococcal
- Pneumococcal
- Polio
- Rotavirus
- Tetanus, diphtheria, pertussis: Tdap preferred; consider an early pertussis booster for high-risk travelers.
- Varicella

Depending on your itinerary, your personal risk factors, and the length of your visit, your health care provider may offer you vaccination against dengue, rabies, or typhoid fever.

Malaria

The following is current information as reported by the World Health Organization (WHO) and the US Centers for Disease Control (CDC):

WHO—International Travel and Health (current online update, Country List)

(2019) Low to moderate malaria risk due predominantly to *P. vivax* exists throughout the year in most of the country. *P. vivax* resistance to chloroquine has been reported. Malaria risk due to *P. falciparum* is still present.

- Recommended prevention: **C** – Risk of *P. falciparum* malaria. Mosquito-bite prevention plus atovaquone-proguanil or doxycycline or mefloquine chemoprophylaxis (select according to drug-resistance pattern, reported side-effects and contraindications).^{a, b}

^aAlternatively, for travel to rural areas with low risk of malaria infection, mosquito bite prevention can be combined with stand-by emergency treatment (SBET).

^bIn certain areas with multidrug-resistant malaria, mefloquine chemoprophylaxis is no longer recommended. At present, these areas include Cambodia, south-eastern Myanmar and Thailand.

WHO Country List footnote: When available, the date of the most recent update or confirmation is indicated in parentheses in the country list. If no accurate date is indicated, the most recent update or confirmation was provided prior to 2013.

CDC—Health Information for International Travel (current online edition)

Transmission areas

- All

Drug resistance¹

- Chloroquine

Species

- *P. vivax* (primarily)
- *P. falciparum* and *P. ovale* (rare)

Recommended chemoprophylaxis

- Atovaquone-proguanil, doxycycline, mefloquine, tafenoquine²

¹ Refers to *P. falciparum* malaria, unless otherwise noted.

² Tafenoquine can cause potentially life-threatening hemolysis in people with glucose-6-phosphate-dehydrogenase (G6PD) deficiency. Test for G6PD deficiency with a quantitative laboratory test before prescribing tafenoquine to patients.

Other Concerns

Travelers' Diarrhea

Moderate risk exists throughout the country, with minimal risk in deluxe accommodations. Community sanitation and food safety measures may be inadequate. Some itineraries (e.g., remote destinations, austere accommodations) and activities (e.g., ecotourism, eating street or local-market food) further increase risk.

Travelers should observe food and beverage precautions, which reduce the likelihood of illness.

Travelers should carry loperamide for self-treatment of diarrhea and, if risk is moderate to high, an antibiotic to add if diarrhea is severe. Consult a knowledgeable health care provider regarding which antibiotic is appropriate for you and most effective for your destination.

Other Food-Borne Illnesses

Precautions to prevent seafood poisoning may be needed.

Other Disease and Health Risks

Additional concerns include helminths, hepatitis C, leptospirosis, marine hazards, sexually transmitted infections, snakebites, tuberculosis.

Safety and Security

Key Safety Risks

- Road conditions
- Domestic air travel
- Maritime safety
- Petty crime
- Heightened crime risk for women
- Natural disasters

Key Security Threats

- Occasional violent crime

Emergency Contacts

The police emergency numbers are 111 and 22222.

© 2026 Shoreland, Inc. All rights reserved.